

## Short term effects of irradiance on the growth of *Pterocladia capillacea* (Gelidiales, Rhodophyta)

RITA F. PATARRA, ANNA A. LLOVERAS, ANA S. CARREIRO, MARIA H. ABREU,  
ALEJANDRO H. BUSCHMANN AND ANA I. NETO



Patarra, R.F., A.A. Lloveras, A.S. Carreiro, M.H. Abreu, A.H. Buschmann and A.I. Neto 2019. Short term effects of irradiance on the growth of *Pterocladia capillacea* (Gelidiales, Rhodophyta). *Arquipelago. Life and Marine Sciences* 36: 85-94. <https://doi.org/10.25752/arq.19682>

*Pterocladia capillacea* has been economically exploited for agar extraction in the Azores for many years. Harvesting dropped to a full stop in the early 1990s due to a population collapse, but restarted in 2013. Since then it has been intensively harvested and overexploitation must be prevented, with both sustainable harvesting and effective cultivation practices. This study represents the first attempt to determine optimal conditions for *P. capillacea* production in the Azores, and evaluates its vegetative growth in two experiments using von Stosch's medium designed to test entire thallus and tips portions response to different irradiances (30, 70 and 150  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). The best relative growth rate (RGR) was recorded at 150  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  for the entire thalli and tips after two-weeks and three-weeks, respectively, indicating that an acclimation period is necessary to assure the growth of this alga under experimental conditions. Higher RGR was obtained at higher irradiance ( $3.98 \pm 2.10\% \text{ fm day}^{-1}$ ), but overall, growth rates were low or negative. Epiphytes were a serious problem towards the end of the entire thallus experiments, where *Feldmannia irregularis* proliferate at all irradiances. Future cultivation approaches complemented with other relevant environmental factors (e.g. pH, photoperiod, salinity), are recommended.

Key words: epiphytes, growth, *in vitro* cultivation, light intensity, *Pterocladia capillacea*

Rita F. Patarra<sup>1,5</sup> (email: [patarrarf@gmail.com](mailto:patarrarf@gmail.com)), A.A. Lloveras<sup>2</sup>, A.S. Carreiro<sup>3</sup>, M.H. Abreu<sup>4</sup>, A.H. Buschmann<sup>5</sup> and A.I. Neto<sup>1</sup>. <sup>1</sup>cE3c - Centre for Ecology, Evolution and Environmental Changes/ Azorean Biodiversity Group and University of Azores, Faculty of Sciences and Technology, Department of Biology, 9501-801 Ponta Delgada, Azores, Portugal. <sup>2</sup>Facultad de Biología, Universidad de Barcelona, Avenida Diagonal, núm. 643, 08028 Barcelona, Spain. <sup>3</sup>University of Azores, Faculty of Sciences and Technology, Department of Biology, 9501-801 Ponta Delgada, Azores, Portugal. <sup>4</sup>ALGAplus, Prod. e Comerc. de Algas e Seus Derivados, Lda. Travessa Alexandre da Conceição, 3830-196 Ílhavo, Portugal. <sup>5</sup>Centro i-mar & CeBiB, Universidad de Los Lagos, Camino Chinquihue km 6, Puerto Montt, Chile.

### INTRODUCTION

*Pterocladia capillacea* (S.G. Gmelin) Santelices & Hommersand is a well-known red alga, recognized to produce a very high quality bacteriological agar and agarose (Bixler & Porse 2010; Boo et al. 2010; Rhein-Knudsen et al. 2015;

Lahaye & Rochas 1991). The demand for agar producing seaweeds is high; nonetheless, all raw materials are obtained from harvests of wild plants, resulting in limited or overexploited natural seaweed stocks (Bixler & Porse 2010; Callaway 2015; Marinho-Soriano 2017). Wild plants of *P. capillacea* are relatively small (up to 17cm), and

mostly inhabit tropical and subtropical waters, although a few extend into temperate latitudes (Lawson & John 1982; Santelices 1998; Patwary et al. 1998; Santelices & Hommersand 1997; Neto 2000a, Neto 2000b; Boo et al. 2010). The growth in length of its axes and branches, as in all genera presently recognized in the Gelidiaceae, depends on the activity of a dome-shaped apical cell (Rodríguez & Santelices 1987; Abbott 1997; Lee 2008). Its erect fronds can persist for up to 7 years although the average is only 2-3 years (Dixon & Irvine 1977). *P. capillacea* growth is site specific and influenced by irradiance, temperature and water movement (e.g. Macler & Zupan 1991; Santelices 1991; Oliveira & Berchez 1993; Friedlander 2008).

Epilithic in the lower intertidal and shallow subtidal (down to 15m), this species is widely distributed in the Azores (Neto 2000b), where it was considerably harvested between 1950-80 t, reaching an annual biomass between 1000-2000 t, (Fralick & Andrade 1981; Santos & Duarte 1991; Melo 2002; Santos & Melo 2018), dried and then exported for agar extraction (Neto et al. 2005). Due to a population collapse in the early 1990s, harvesting dropped to a full stop, but restarted in 2013 and, at the present, *P. capillacea* is being harvested in six of the nine Islands of the Archipelago (Terceira, São Miguel, Graciosa, Faial, São Jorge and Pico Islands; LOTAÇOR 2019) and sold almost exclusively to IBERAGAR S.A. and TAE Lda. companies (AOnline 2016; unpublished observations). In 2016, 450 wet weight (ww) tonnes of *P. capillacea* were harvested, representing a total value of 450000 € (GaCS 2017), corresponding grossly to 1 € Kg<sup>-1</sup> ww. Due to this recent resurgence of the exploitation, there is an urgent need to implement good management practices for the harvest of the wild resource and to promote the species cultivation (Patarra et al. 2014; Rebours et al. 2014; Netalgae 2012; Patarra 2018).

Today there is not any known commercial cultivation of this Gelidiales (Melo 1998; Friedlander 2008; Callaway 2015; Marinho-Soriano 2017; Santos & Melo 2018), mostly due to its slow growth rates (Friedlander & Lipkin 1982; Friedlander & Zelikovitch 1984; Macler & Zupan 1991; Yokoya & Oliveira 1992; Oliveira &

Berchez 1993; Felicini et al. 2002; Fujimoto et al. 2014). However, land base studies on *P. capillacea* in Israel, revealed promising growth rates per week (28.3% in average) during winter time (Gal-Or & Israel 2004). According to a revision of the Gelidiales cultivation made by Friedlander (2008) distinct types of seaweed material, namely different sized portions from different parts of the thallus can have different growth. This was earlier observed by Rodríguez (1996) who reported that the vegetative growth in length and branch proliferation of *Gelidium sclerophyllum* was favored by medial fragments, whereas the rhizoidal propagation was favoured by apices.

To our best knowledge, a single study on the physiological responses of Azorean Gelidiales was performed (Fralick et al. 1990), and authors indicated that optimal photosynthesis activity occurred at 177  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , between 15 and 25°C. To move to an effective culture of this species, a clarification of the most relevant factors and constraints for its production is needed. The present work is a first attempt to understand how different irradiances affect the vegetative growth of distinct portions of the thallus (entire thallus and tips) of *P. capillacea* from the Azores.

## MATERIAL AND METHODS

### Experimental setup

Sterilized natural seawater (SNW) was prepared by filtering the water with 0.2  $\mu\text{m}$  pore size filter and autoclaved. Von Stosch's enrichment culture medium (VSE) was prepared according to Guiry & Cunningham (1984). All the material used during media preparation and in the experiments was sterilized either by autoclave (2 atm, 120°C, 20 min) or by using a drying oven (150°C, 2 hr).

FronDS were collected by hand, between December 2013 and February 2014, from the low intertidal zone of a rocky shore of the south coast of São Miguel Island (37° 50' N, 25° 30' W; Azores archipelago), where stable populations of *P. capillacea* are known to occur (Neto 2000b, Neto 2000a, Neto 2001, Wallenstein et al. 2008). FronDS were transported to the laboratory within 4 hr of the collection and kept in natural seawater tanks (5 L). At each collection time, voucher

specimens were made and deposited in the Herbarium Ruy Telles Palhinha (AZB) of the Biology Department, University of the Azores (*P. capillacea*, AZB- SMG-13-90, AZB-SMG-14-01, AZB-SMG-14-07).

One week before the experiments, algae were incubated in 1 L Erlenmeyer flasks containing 500 ml of SNW enriched with VSE and kept in constant movement through bottom aeration.

A preliminary experiment was conducted to evaluate the effect of different pre-cleaning solutions (sodium hypochlorite 1%; betadine 10% and SNW) on the growth of epiphytes and endophytes of *P. capillacea* (Table 1). The pre-cleaning solution Bleach promoted the growth of the red endophytes *Anotrichium* spp. and *Epicladia* spp. (unpublished data). Based on these preliminary results, the pre-cleaning solution Betadine 10% was used in all subsequent experiments. For this, fronds of *P. capillacea* were rinsed under tap water and visible epibionts and surface sediments were removed by hand and gently brushed if necessary. Then, algae were checked under a dissecting microscope and the healthy ones retained for the cultivation essays. Entire thalli (of 2-3cm long) and tips portions (1cm long) were cleaned in SNW, submerged for 30s in the pre-cleaning solution of Betadine 10% and cleaned again in SNW; then they were randomly assigned to experimental units, 1 L Erlenmeyer flasks each containing eight individual entire thallus or tips portions (depending on the subsequent experiment) and 500 ml of SNW enriched with VSE. The growth of excised fragments was evaluated in two experiments designed to test if distinct algae material: i) entire thallus (ET) and ii) tips portions (TP) respond differently under different photon flux densities (PFD, 30, 70 and 150  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ).

The experiments were carried in cultivation chambers (Sanyo MLR-351; Japan) using a photoperiod of 12L:12D and 18°C temperature as fixed factors and lasted 5 weeks. The temperature of 18°C was used to match natural conditions of the collecting site and the average annual sea surface temperature (SST) in the Azores archipelago (DETRA 2013, Patarra *et al.* 2017). The 12L:12D photoperiod chosen in the experiments relies on the fact that there is no

consensus on the best L:D combination for *P. capillacea* growth (e.g. 10:14 LD, Fralick *et al.* 1990, 12:12 L:D; Nasr *et al.* 1966, 16.8 LD; Yokoya & Oliveira 1992, 12:12 LD; Felicini *et al.* 2002). The irradiance was measured with a LI-250 Light Meter (LI-COR, USA) inside the culture chambers and whenever the algae were collected in the field. When the culture medium was changed, the algal material was blotted dry and weighted as fresh mass (fm) using a digital balance (Kern ALJ220-5DNM, Germany;  $\pm 0.0001$  g precision). In each of these occasions, algae were placed on plasticized graph paper and photographed (Sony  $\alpha 230$  with an objective of 18-55mm, Japan) for posterior length analyses. The pH was monitored (three measurements per replicate) with a pH sensor (Hanna Instruments, HI98127 USA).

### Growth responses

Once a week, fresh biomass (g) and length (cm) were registered, and relative growth rate was calculated (RGR; % fm day<sup>-1</sup>) for each replicate (n=4, SE) in all treatments. Length of individual thallus/portions was obtained from the above mentioned images using the AxioVision LE Digital Imaging Software (SE 64 Rel. 4.9.1., Germany) and by measuring each individual thallus/portion from its base to the top. At each sampling time the RGR was calculated for each replicate flask according to the following formula (Abreu *et al.* 2011):

$$\text{RGR} = \frac{[\text{Ln}(\text{final fm}) - \text{Ln}(\text{initial fm})]}{\text{time (days)} \times 100}$$

These calculations were based on the sum value of the eight individual thallus/portions in each flask. For the analysis (see section 2.3), the mean value of weekly growth rates measured throughout the experiment was used to account for temporal growth rates variation.

### Data analysis

For all analysis, we used flasks as sampling units. One-way analysis of variance (ANOVA, e.g. Underwood 1997; PFD as a fixed factor with 3 levels, 4 replicates each) was used to test for differences in RGR in the different treatments.

Prior to the analysis, Cochran's test was used to test for homogeneity of variances and transformations were applied whenever necessary. When this was not possible, analyses were run on the untransformed data, since ANOVA is robust to departures from this assumption when designs are balanced and replication is high (Underwood 1997). However, a more conservative p-value ( $\alpha = 0.01$ ) was used. The *a posteriori* Student–Newman-Keuls (SNK) test was used to investigate differences among levels within significant terms. All statistical analyses were performed using the software

GMAV v5.

## RESULTS

The abiotic factors (temperature, irradiance) monitored during the experiment were kept in the pre-defined range. Carbon availability was always in optimal levels, as determined by the pH recorded values, ranging between  $7.00 \pm 0.01$  and  $7.93 \pm 0.02$  in ET experiment and  $7.02 \pm 0.05$  and  $7.90 \pm 0.02$  in TP experiment.

Table 1. List of epiphytes and endophytes identified at the end of the preliminary experiment. SNW - sterilized natural seawater.

Taxa	Authorities	Occurrence	Pre-cleaning solutions		
			SNW	Betadine 10%	Bleach
<b>RHODOPHYTA</b>					
<i>Anotrichium tenue</i>	Nägeli	endophyte			x
<i>Bangia</i> spp.	Lyngbye	epiphyte			x
<i>Ceramium rubrum</i>	C. Agardh	epiphyte			x
<b>CLOROPHYTA</b>					
<i>Cladophora</i> spp.	Kützing	epiphyte		x	
<i>Enteromorpha multiramosa</i>	Bliding	epiphyte	x		
<i>Epicladia</i> spp.	Reinke	endophyte			x
<i>Ulva compressa</i>	Linnaeus	epiphyte	x		x
<i>Ulva rigida</i>	C. Agardh	epiphyte		x	x
<b>HETEROKONTOPHYTA</b>					
<i>Feldmannia irregularis</i>	(Kützing) Hamel	epiphyte	x	x	

In the ET experiment there was no biomass increase in any of the treatments for the first 3 weeks (Figure 1A, B). However, pronounced weight and length increments were recorded after the third week for the  $150 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  treatment (Figure 1A), and the same pattern occurred from the fourth to the fifth week at  $70 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Figure 1A). However, in both situations an increase in the epiphytes (mostly *Feldmannia irregularis*) growth (from the base to the top of the thalli) was recorded, as well as an accumulation of a layer of died cells over the thalli surface. Due to the amount of epiphytes growing

at  $150 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  at the fifth week, no measurements were taken from that moment forward and for RGR statistical analysis only 30 and  $70 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  treatments were used (Table 2). The lower RGR was observed at  $30 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  ( $-2.02 \pm 1.25\% \text{ fm day}^{-1}$ ) indicating loss of tissue (Figure 1C, Table 2). Overall, the best entire thalli RGR was recorded at  $150 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  ( $3.98 \pm 2.10\% \text{ fm day}^{-1}$ , Figure 1C).

As for the TP experiment, there was a similar growth pattern for the  $70$  and  $150 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  treatments, with slight mass increments

Short term effects of irradiance on the growth of *Pterocliadiella capillacea*




Fig. 1. Effect of photon flux density (PFD) on the growth of *P. capillacea* entire thallus (ET; panels A, B and C) and tips portions (TP; panels D, E and F) cultivated at three PFD (30, 70 and 150  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ), 18 °C and 12:12 L:D photoperiod, after five weeks in culture. Fresh mass and length (mean  $\pm$  SE, n=4); relative growth rate (RGR, mean  $\pm$  SE, n=20;  $p < 0.05$ ). # mean RGR of four weeks, not used in the statistical analysis.

after two weeks in culture (Figure 1D, E). No recovery was observed but the growth was lower at the higher irradiance (150  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) (Figure 1E, Table 2) and no epiphytes were recorded. Nevertheless, RGR was negative at all

PFD (Figure 1F). As for the ET, the best growth was obtained for the higher PFD used. Experiments lasted 5 weeks, since then algae were mushy and colorless (some of them even white).

Table 2. Univariate analysis of variance (ANOVA) examining the effect of photon flux density (two levels for entire thallus experiment, ET; three levels for tips portions experiment, TP; fixed) on the relative growth rate of *P. capillacea* cultivated at 18 °C, and 12:12 L:D photoperiod, after 5 weeks in culture.

Source	ET				TP				
	df	MS	F	P	df	MS	F	P	
PFD	1	86.44	2.42	0.1280	2	6.70	0.21	0.8135	
Residual	38	35.71			57	32.36			
Cochran's test		C = 0.67 ns					C = 0.47 ns		
Transformation		none					none		

PFD = photon flux density; ET = entire thallus experiment; TP = tips portions experiment; n.s., not significant

## DISCUSSION

The best growth performance observed at high PFD for both, the entire thallus and the tip portions, submitted to an longer acclimation period (seen as necessary to assure the growth of *P. capillacea* under experimental conditions) is in agreement with reports for this species by other researchers (e.g. Nasr et al. 1966; Correa et al. 1999; Felicini et al. 2002). However, the unexpected decrease in length observed for both entire thallus and plant tips is in disagreement with reports by other authors, which reported that, although *P. capillacea* exhibited low growth rates, it maintained a viable growth for several days or weeks in artificial conditions (e.g. Stewart 1984; Macler & Zupan 1991; Yokoya & Oliveira 1992; Oliveira & Berchez 1993, Felicini et al. 2002; Friedlander 2008; Fujimoto et al. 2014). This may be related with the collection time of the algae material used to run laboratory experiments. In fact, Stewart (1984) reported a best initial growth and survival for cultures initiated with Californian of *P. capillacea* collected between October and March, corresponding to local late autumn/winter period. Similar results were reported by Gal-Or & Israel (2004) for Israeli material. In the present study all the laboratory trials were run at 18 °C (the mean annual SST in the Azores Archipelago, DETRA 2013, Patarra et al. 2017), but the algae were collected from December to early Spring when SST varies between 16 and 17 °C (DETRA 2013). It is likely that the observed negative growth could be related to higher temperature used. In fact, a poor growth performance of *P. capillacea* at high temperatures has been reported by several authors (Fralick et al. 1990; Yokoya & Oliveira 1992; Gal-Or & Israel 2004; Fujimoto et al. 2014). It is, however, worth considering, that *P. capillacea* in the Azores is distributed from low intertidal to shallow subtidal zones (Neto 2000a, Neto 2000b, Neto 2001; Wallenstein et al. 2008), being adapted to a wide range of irradiances and water temperatures, although the temperature range in shallow subtidal zones does not vary greatly (HI 2000). Furthermore, Fralick & Andrade (1981) observed that wild Azorean plants exhibited optimal growth at 20-22 °C, coinciding

to the period between June and October when the irradiance level is higher in the Azores.

Regarding irradiance, the obtained mean growth rates were negative/low at all PFD, and, even in the higher irradiances, their was loss of tissue, and the thallus was mushy, colorless or even white at the end of the experiments. This is in disagreement with reports by other researchers for Gelidiales: Macler & West (1987) kept vegetative cultures of *Gelidium coulteri* for more than a year with an exponential growth rate less than 10% day<sup>-1</sup> at 150  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ; Fralick et al. (1990) in a research on the physiological responses of *Pterocladia* and *Gelidium* (Gelidiales, Rhodophyta) from the Azores reported optimal photosynthesis for *P. capillacea* at 177  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  with oxygen production values remaining high until 320  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ; Rico (1991) reported maximum growth rates (10.0% day<sup>-1</sup>) for *Gelidium pulchellum* at 130- $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ; Sousa-Pinto et al. (1999) observed that the growth of *Gelidium pulchellum* pieces increased with irradiance up to 130  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ; Gal-Or & Israel (2004) recorded maximal growth rates for *P. capillacea*, using a gradient table, at 100  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  with winter plants; Harb et al. (2018) reported a higher growth rate for *P. capillacea* at 300  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ , as long as increasing photosynthetic pigment and protein contents.

Worth considering, however, that Stewart (1984) reported that higher levels of irradiance favored contaminants and were correlated with larger numbers of bleached *P. capillacea* thalli that did not exhibit measurable growth. This is similar to what we observed in the present study, in which the growth of the epiphyte *Feldmannia irregularis* at 150  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  after 4 weeks in culture was a serious problem (growing from the base to the top of the thalli). According to Friedlander (2008) one of the biological factors that most affects *Gelidium* cultivation is indeed the growth of epiphytes. Sousa-Pinto et al. (1999) in their study on the effect of light on the growth of *Gelidium pulchellum* reported that the algae were heavily epiphytized (with cyanobacteria, fungi and algae) at the higher irradiances (240-430  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). Fungi and cyanobacteria

proliferation were inhibited with antibiotics, but green algae persisted and were only marginally contained after washing the thalli with distilled water in-between medium changes. Buschmann et al. (1997) also reported problems with Ceramialean epiphytes on *Gracilaria* and observed that their abundance increased significantly from the apical (new tissues) to the central parts of the thalli (older tissues). Probably the lower load of epiphytes observed on the apical tips portions of *P. capillacea* in the present study is related to the new age of this tissue, but could be also related with the adopted cleaning procedures. From the known cleaning methods (e.g.: i) cleaning the algae with a soft brush; ii) wash them with distilled water; iii) using pre-cleaning solutions, such as sodium hypochlorite, Betadine 10%, antibiotics) used by several authors (see Salinas 1991; Sousa-Pinto et al. 1999; Redmond et al. 2014), the one adopted in the present study (pre cleaning followed by the use of Betadine 10%) seemed to be effective for some epiphytes identified in the preliminary experiments with the tip portions (TP).

The obtained values for the culture medium pH in all experiments suggest the algae were not CO<sub>2</sub> limited (in accordance to Lignell & Pedersén 1989; Hargreaves 1998; Mercado et al. 2001).

Altogether, the adopted combination of irradiance, temperature, nutrients and water aeration in the present study was not the best option for the growth of *P. capillacea* from the Azores. Nonetheless, small studies, as the present ones, are an important tool for the collection of data on the biology of the species, that could be used in future long-term experiments. Considering the economic importance of the species in the region, future cultivation approaches using different culture combinations, complemented with relevant environmental factors (e.g. pH, photoperiod, salinity) and a delicate selection of fast growing ecotypes are recommended.

#### ACKNOWLEDGEMENTS

This study was partially supported by project AQUAIMPROV, ON.2 – O Novo Norte, (SAESCTN-PII&DT/1/2011); Portuguese National

Funds, through FCT – Fundação para a Ciência e Tecnologia, within the projects UID/BIA/00329/2013, 2015 - 2018 and UID/BIA/00329/2019, CIRN (Centro de Investigação de Recursos Naturais, University of the Azores), and CIIMAR (Interdisciplinary Centre of Marine and Environmental Research, Porto, Portugal). RFP was supported by a doctoral grant M3.1.2/F/024/2011, Fundo Regional para a Ciência e Tecnologia. The authors are grateful to Pedro Pereira and Natália Cabral (Department of Biology, Azores University) for their technical support during the experimental studies. We would like to thanks Dr. Eva Cacabelos for her support on the seawater collection along the experimental period. Special thanks go to Dr. Gustavo Martins for his helpful comments on the statistical analysis.

#### REFERENCES

- Abbott, I. A. 1997. *Taxonomy of economic seaweeds: With reference to some pacific species*, Volume VI. University of California, La Jolla, California. 212 pp.
- Abreu, M. H., Pereira, R., Yarish, C., Buschmann, A. H. and I. Sousa-Pinto 2011. IMTA with *Gracilaria vermiculophylla*: Productivity and nutrient removal performance of the seaweed in a land-based pilot scale system. *Aquaculture* 312:77-87.
- AOnline 2016. Apanha de algas rendeu 120 mil euros em 2015. Seaweed harvest has yielded 120000 € in 2015. *Açoriana Oriental* Ponta Delgada. Available from: <http://www.acorianooriental.pt/pagina/edicao-impressa/2016-02-21/regional> (cited 21 February 2016).
- Bixler, H. J. and H. Porse 2010. A decade of change in the seaweed hydrocolloids industry. *Journal of Applied Phycology* 23:321-335.
- Boo, S. M., Kim, S. Y., Hong, I. S. and I.K. Hwan 2010. Reexamination of the genus *Pterocladia* (Gelidiaceae, Rhodophyta) in Korea based on morphology and rbcL sequences. *Algae* 25:1-9.
- Buschmann, A. H., Retamales, C. A. and C. Figueroa 1997. Ceramialean epiphytism in an intertidal *Gracilaria chilensis* (Rhodophyta) bed in southern Chile. *Journal of Applied Phycology* 9:129-135.
- Callaway, E. 2015) Lab staple agar runs low. *Nature* 528:171-172.
- Correa, J. A., Beltrán, J., Buschmann, A. H. and R.

- Westermeier 1999. Healing and regeneration responses in *Gigartina skottsbergii* (Rhodophyta, Gigartinales): Optimization of vegetative propagation for cultivation. *Journal of Applied Phycology* 11:315-327.
- DETRA 2013. Implementation of Remote Sensing Techniques in the Azores – DETRA project. Sea surface temperature 1997-2010. DOP/UAç.
- Dixon, P. S. and L.M. Irvine 1977. *Seaweeds of the British Isles*. Volume 1. Rhodophyta. Part 1. Introduction, Nemaliales, Gigartinales, British Museum (Natural History), London. 252 pp.
- Felicini, G. P., Bottalico, A. and M. Fanelli 2002. Morphogenesis in *Pterocladia capillacea* (Rhodophyta, Gelidiales): bud differentiation in relation to irradiance-temperature combinations. *Plant Biosystems* 136:261-267.
- Fralick, R. A. And F. Andrade 1981. The growth, reproduction, harvesting and management of *Pterocladia pinnata* (Rhodophyceae) in the Azores, Portugal. Pp. 289-295 in: T. Levring (Ed.). Proceedings of the Tenth International Seaweed Symposium, Gothenburg (Sweden), 1980. Walter de Gruyter and Co., Gothenburg, Sweden.
- Fralick, R. A., Baldwin, H. P., Neto, A. I. and E.J. Hehre 1990. Physiological responses of *Pterocladia* and *Gelidium* (Gelidiales, Rhodophyta) from the Azores, Portugal. *Hydrobiologia* 204-205:479-482.
- Friedlander, M. (2008). Advances in cultivation of Gelidiales. *Journal of Applied Phycology* 20:451-456.
- Friedlander, M. and Y. Lipkin 1982. Rearing of agarophytes and carrageenophytes under field conditions in the Eastern Mediterranean. *Botanica Marina* 25:101-105.
- Friedlander, M. and N. Zelikovitch 1984. Growth rates, phycocolloid yield and quality of the red seaweeds, *Gracilaria* sp., *Pterocladia capillacea*, *Hypnea musciformis*, and *Hypnea cornuta*, in field studies in Israel. *Aquaculture* 40:57-66.
- Fujimoto, M., Nishihara, G. N. and R. Terada 2014. The effect of irradiance and temperature on the photosynthesis of two agarophytes *Gelidium elegans* and *Pterocladia tenuis* (Gelidiales) from Kagoshima. *Japan. Fisheries Science* 80:695-703.
- GaCS 2017. Governo Regional estimula atividades que geram rendimentos complementares aos pescadores açorianos. Presidência do Governo Regional dos Açores. Gabinete de Apoio à Comunicação Regional. Available from: <http://www.azores.gov.pt/GaCS/Noticias/2017/Junho/Governo+Regional+estimula+atividades+que+geram+rendimentos+complementares+aos+pescadores+a%C3%A7orianos.htm> (cited 15 June 2017).
- Gal-Or, S. and A. Israel. 2004. Growth responses of *Pterocladia capillacea* (Rhodophyta) in laboratory and outdoor cultivation. *Journal of Applied Phycology* 16:195-202.
- Guiry, M. D. and E.M. Cunningham 1984. Photoperiodic and temperature responses in the reproduction of north-eastern Atlantic *Gigartina acicularis* (Rhodophyta: Gigartinales). *Phycologia* 23:357-367.
- Harb, T. B., Nardelli, A. and F. Chow 2018. Physiological responses of *Pterocladia capillacea* (Rhodophyta, Gelidiales) under two light intensities. *Photosynthetica* 56(4):1093 - 1106
- Hargreaves, J. A. 1998. Nitrogen biogeochemistry of aquaculture ponds. *Aquaculture* 166:181-212.
- Instituto Hidrográfico 2000. *Roteiro da Costa de Portugal - Arquipélago dos Açores*. Instituto Hidrográfico, Lisboa.
- Lahaye, M. and C. Rochas 1991. Chemical structure and physico-chemical properties of agar. *Hydrobiologia* 221:137-148.
- Lawson, G. W. and D.M. John 1982. The marine algae and coastal environment of Tropical West Africa. *Pterocladia* J. Agardh 1852, *Nova Hedwigia*, 70. J. Cramer, Vaduz. 455 pp.
- Lee, R. E. 2008. *Phycology*. Cambridge University Press, Cambridge. 547 pp.
- Lignell, Å. And M. Pedersen 1989. Effects of pH and inorganic carbon concentration on growth of *Gracilaria secundata*. *British Phycological Journal* 24:83-89.
- LOTAÇOR 2019. Total annual yields (landings in tonnes) of *Pterocladia capillacea* harvested in the Archipelago of the Azores. Lotaçor S.A. - Serviço de Lotas dos Açores. Available from: [http://www.lotacor.pt/PescadoDescarregadoAuth/g\\_ratis.php](http://www.lotacor.pt/PescadoDescarregadoAuth/g_ratis.php). (cited 14 September 2019).
- Macler, B. A. and West, J. A. 1987. Life history and physiology of the red alga, *Gelidium coulteri*, in unialgal culture. *Aquaculture* 61:281-293.
- Macler, B. A. and J.R. Zupan 1991. Physiological basis for the cultivation of the Gelidiaceae. *Hydrobiologia* 221:83-90.
- Marinho-Soriano, E. 2017. Historical context of commercial exploitation of seaweeds in Brazil. *Journal of Applied Phycology* 29:665-671.
- Melo, R. A. 1998. Gelidium commercial exploitation: natural resources and cultivation. *Journal of Applied Phycology* 10:303-314.
- Melo, R. A. 2002. Exploração dos recursos algológicos em Portugal. Pp. 45-65 in: M. A. C. Martins-Loução (Ed.). *Fragmentos em Ecologia*. FCUL-Livraria Escolar Editora. 355 pp.
- Mercado, J. M., Niell, F. X. and M.C. Gil-Rodríguez 2001. Photosynthesis might be limited by light, not inorganic carbon availability, in three intertidal

- Gelidiales species. *New Phytologist* 149:431-439.
- Nasr, A. H., Mohsen, A. F. and I.A. Bekheet 1966. Effect of salinity and temperature variations on *Pterocladia capillacea*. *Hydrobiologia* 27:395-400.
- Netalgae 2012. Seaweed industry in Europe: a guide to better practice. An output of the netalgae project. Available from: [http://www.netalgae.eu/uploadedfiles/Guide\\_8p\\_UK.pdf](http://www.netalgae.eu/uploadedfiles/Guide_8p_UK.pdf) (cited April 2019)
- Neto, A. I. 2000a. Ecology and dynamics of two intertidal algal communities on the littoral of the island of São Miguel (Azores). *Hydrobiologia* 432:135-147.
- Neto, A. I. 2000b. Observations on the biology and ecology of selected macroalgae from the littoral of São Miguel (Azores). *Botanica Marina* 43:483-498.
- Neto, A. I. 2001. Macroalgal species diversity and biomass of subtidal communities of subtidal communities of São Miguel (Azores). *Helgoland marine research* 55:101-111.
- Neto, A. I., Tittley, I. and P. Raposeiro 2005. Flora Marinha do Litoral dos Açores. Rocky shore marine flora of the Azores. Secretaria Regional do Ambiente e do Mar, Horta, Portugal. 156 pp.
- Oliveira, E. C. and F.A.S. Berchez 1993. Resource biology of *Pterocladia capillacea* (Gelidiales, Rhodophyta) populations in Brazil. *Hydrobiologia* 260-261:255-261.
- Patarra, R. 2018. Culture studies of economically important seaweeds. Pages viii+157 pp. +Appendix. Faculdade de Ciências e Tecnologia, Departamento de Biologia. Doctoral thesis in Biology, University of the Azores, Faculty of Sciences and Technology, Department of Biology, Ponta Delgada.
- Patarra, R. F., Buschmann, A. H., Abreu, M. H. and A. I. Neto 2014. Seaweed cultivation in the Azores ... Opportunities and challenges. Cultivo de macroalgas nos Açores... Oportunidades e desafios. *Boletim de Biotecnologia da Sociedade Portuguesa de Biotecnologia* 2:16-18.
- Patarra, R. F., Carreiro, A. S., Lloveras, A. A., Abreu, M. H., Buschmann, A. H. and A.I. Neto 2017. Effects of light, temperature and stocking density on *Halopteris scoparia* growth. *Journal of Applied Phycology* 29:405-411.
- Patwary, M. U., Sensen, C. W., MacKay, R. M. and J. P. van der Meer 1998. Nucleotide sequences of small-subunit and internal transcribed spacer regions of nuclear rRNA genes support the autonomy of some genera of the Gelidiales (Rhodophyta). *Journal of Phycology* 34:299-305.
- Rebours, C., Marinho-Soriano, E., Zertuche-González, J. A., Hayashi, L., Vásquez, J. A., Kradolfer, P., Soriano, G., Ugarte, R., Abreu, M. H., Bay-Larsen, I., Hovelsrud, G., Rødven, R. and D. Robledo 2014. Seaweeds: an opportunity for wealth and sustainable livelihood for coastal communities. *Journal of Applied Phycology* 26:1939-1951.
- Redmond, S., Green, L., Yarish, C., Kim, J. and C. Neefus 2014. *New England seaweed culture handbook* - Nursery systems, Connecticut Sea Grant CTSG-14-01. 92 pp.
- Rhein-Knudsen, N., Ale, M. T. and A.S. Meyer, A. S. 2015. Seaweed hydrocolloid production: an update on enzyme assisted extraction and modification technologies. *Mar Drugs* 13:3340-3359.
- Rico, J. M. 1991. Field studies and growth experiments on *Gelidium latifolium* from Asturias (northern Spain). *Hydrobiologia* 221:67-75.
- Rodríguez, D. 1996. Vegetative propagation by fragmentation of *Gelidium sclerophyllum* (Gelidiales, Rhodophyta). *Hydrobiologia* 326:361-365.
- Rodríguez, D. and B. Santelices 1987. Patterns of apical structure in the genera *Gelidium* and *Pterocladia* (Gelidiaceae, Rhodophyta). *Hydrobiologia* 151:199-203.
- Salinas, J. M. 1991. Spray system for re-attachment of *Gelidium sesquipedale* (Clem.) Born. et Thur. (Gelidiales: Rhodophyta). *Hydrobiologia* 221:107-117.
- Santelices, B. 1991. Production ecology of *Gelidium*. *Hydrobiologia* 221: 31-44.
- Santelices, B. 1998. Taxonomic review of the species of *Pterocladia* (Gelidiales, Rhodophyta). *Journal of Applied Phycology* 10:237-251.
- Santelices, B. and M. Hommersand 1997. *Pterocladia*, a new genus in the Gelidiaceae (Gelidiales, Rhodophyta). *Phycologia* 36:114-119.
- Santos, R. and P. Duarte 1991. Marine plant harvest in Portugal. *Journal of Applied Phycology* 3:11-18.
- Santos, R. and Melo, R.A. 2018. Global shortage of technical agars: back to basics (resource management). *Journal of Applied Phycology* 30: 2463-2473.
- Sousa-Pinto, I., Murano, E., Coelho, S., Felga, A. and R. Pereira 1999. The effect of light on growth and agar content of *Gelidium pulchellum* (Gelidiaceae, Rhodophyta) in culture. *Hydrobiologia* 398-399: 329-338.
- Stewart, J. G. 1984. Vegetative growth rates of *Pterocladia capillacea* (Gelidiaceae, Rhodophyta). *Botanica Marina* 27:85-94.
- Underwood, A. 1997. *Experiments in ecology: their logical design and interpretation using analysis of variance*. Cambridge University Press, Cambridge. 524 pp.
- Wallenstein, F. F. M. M., Neto, A. I., Álvaro, N. V. and I. Tittley 2008. Subtidal rocky shore communities of the Azores: Developing a biotope survey method.

*Journal of Coastal Research* 1:244-249.  
Yokoya, N. S. and E.C. Oliveira 1992. Temperature responses of economically important red algae and their potential for mariculture in Brazilian waters. *Journal of Applied Phycology* 4:339-345.

*Received 02 Oct 2019. Accepted 22 Oct 2019.  
Published online 18 Nov 2019.*