

Antioxidant activity of macroalgae from the Azores

LISETE S. PAIVA, R.F. PATARRA, A.I. NETO, E.M.C. LIMA & J.A.B. BAPTISTA



Paiva, L.S., R.F. Patarra, A.I. Neto, E.M.C. Lima & J.A.B. Baptista 2012. Antioxidant activity of macroalgae from the Azores. *Arquipelago. Life and Marine Sciences* 29: 01-06.

Free radical scavenging activity (FRSA) was studied to determine the antioxidant activity of 8 common macroalgae species found in the Azores. Species under study were *Porphyra* sp., *Osmundea pinnatifida*, *Pterocliadiella capillacea*, *Sphaerococcus coronopifolius* and *Gelidium microdon*, from Rhodophyta; *Ulva compressa* and *Chaetomorpha pachynema*, from Chlorophyta and *Fucus spiralis* from Phaeophyta. The antioxidant activity was evaluated in methanolic extract by a decolourisation solution test of 2,2-diphenyl-1-picrylhydrazyl (DPPH), used as a stable radical. The methanolic extracts were obtained from dried biomass by sequential extractions, attaining a final concentration of 2 mg.mL⁻¹. The FRSA values ranged from 19.54% for *S. coronopifolius* to 60.05% for *F. spiralis* with standard deviation (SD) varying between 1.41% and 6.80%. Results indicated that the studied seaweeds are a very promising source of biological active compounds with antioxidant properties. The seaweeds were collected in the Azorean islands where seawater pollution levels are low. Consequently, these seaweeds represent a valuable and good source of antioxidant material with superior beneficial effects on human health.

Key words: DPPH, functional food, macroalgae screening, radical scavenging activity

Lisete S. Paiva¹ (e-mail: lisetepaiva@gmail.com), Elisabete M.C. Lima² & José A.B. Baptista³, Departamento Ciências Tecnológicas e Desenvolvimento (DCTD), Universidade dos Açores, Rua da Mãe de Deus, 9501-801 Ponta Delgada, Açores, Portugal; ¹²³ & Rita F. Patarra, Centro de Investigação de Recursos Naturais (CIRN), Universidade dos Açores, Rua da Mãe de Deus, 9501-801 Ponta Delgada, Açores, Portugal; Rita F. Patarra & Ana I. Neto, Secção Biologia Marinha, Laboratório Ficologia, Departamento Biologia, Universidade dos Açores, Rua da Mãe de Deus, 9501-801 Ponta Delgada, Açores, Portugal; & Centro Interdisciplinar de Investigação Marinha e Ambiental (CIIMAR), Rua dos Bragas 289, 4050-123 Porto, Portugal.

INTRODUCTION

In living organisms, reactive oxygen species (ROS) and reactive nitrogen species (RNS) are known to cause damage to biomolecules, such as lipids, nucleic acids, proteins, enzymes and other cellular constituents (Halliwell 1991). The most important reaction is lipid peroxidation of unsaturated fatty acids. This oxidative degradation of lipids, proceeds by a free radical chain reaction mechanism, has been known to cause many pathological effects (Spencer et al. 1994). A lipid radical (L•) from a lipid (LH) is formed in the initiation step that then reacts with oxygen to

form a lipid peroxy radical (LOO•) which reacts with an additional lipid molecule to give a lipid hydroperoxide (LOOH) in the propagation step. Fatty acid hydroperoxides are known to be one of the active oxygen species (Ohkawa et al. 1979). The degradation and/or modification of these molecules have been related with various chronic diseases implicated in the processes of aging, as well as in a wide range of degenerative diseases, such as: coronary heart disease, atherosclerosis, cancer, cataracts, diabetes, liver injury, Alzheimer's and Parkinson's diseases, muscular dystrophy and some others neurological disorders (Duan et al. 2006). It is well known that antioxi-

dants can neutralise potentially harmful reactive free radicals in body cells before they cause lipid damage. In addition, lipid peroxidation is one of the major causes of deterioration during storage and food processing. To compensate these deleterious effects, synthetic antioxidant compounds, such as butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) are commonly used, which in excess may contribute to some negative health side effects, according to the published literature (Santoso et al. 2004). Therefore, there is a growing interest on the discovery of natural antioxidants, mainly for two reasons: (1) epidemical and clinical evidence suggests that consumption of fruits and vegetables reduce the risk of developing chronic disease, e.g. cancer and coronary heart disorders, and (2) phytochemicals are generally safer than synthetic chemicals (Dastmalchi et al. 2007). For these reasons, many products with antioxidant properties are widely used, particularly from natural sources, in order to minimise oxidative damage to living cells and to prevent oxidative deterioration (Rekka & Kourounakis 1991; Conforti et al. 2005; Rackova et al. 2007). Antioxidant activity of seaweeds is being intensively investigated due to the currently growing demand from the food and pharmaceutical industries where there is interest in anti-aging and anticarcinogenic natural bioactive compounds (Matsukawa et al., 1997; Siriwardhana et al. 2003; Dias et al. 2005; Senevirathne et al. 2006; Kuda et al. 2005; Ganesan et al., 2008). Furthermore, seaweeds have also been used as a source of novel functional foods with potential nutritional benefits (Santoso et al. 2004). More recently, much attention has been paid to the anti-tumour and anticholesterolemic activities of seaweed constituents (Park et al. 2005; Villaño et al. 2007; Duh 1998). Like plants, seaweeds contain various types of inorganic (high levels of minerals) and organic compounds (vitamins, proteins, essential amino acids, indigestible carbohydrates and dietary fibre) (Jiménez-Escrig et al. 2001; Yoshie et al. 2002; Patarra et al. 2011).

The objective of our study was to determine the antioxidant activity of the most common macroalgae in the Azores islands taking into account the low pollution levels of seawater in the region (Neto et al. 2009) and, consequently, the great

potential of its use for human consumption and/or for the extraction of novel compounds with added value for the food and pharmaceutical industries. In addition, this study also comparatively evaluated the FRSA of the referred macrophytes with synthetic antioxidant BHT.

MATERIAL AND METHODS

SEAWEEDS SAMPLES AND CHEMICALS

All seaweed samples used in this study were collected during January/February 2007, April 2008 and June 2009, from the littoral zone of São Miguel Island, in the Azores Archipelago. The seaweeds were washed with water and alcohol, air-dried and stored in an air-tight container in a freezer (-20 °C) until further analysis. Seaweeds analysed belonged to Phaeophyta (*Fucus spiralis* Linnaeus), Chlorophyta (*Ulva compressa* Linnaeus and *Chaetomorpha pachynema* (Montagne) Kützinger), and Rhodophyta (*Porphyra* sp. C. Agardh, *Osmundea pinnatifida* (Hudson) Stackhouse, *Pterocladia capillacea* (S.G. Gmelin) Santelices & Hommersand, *Sphaerococcus coronopifolius* Stackhouse and *Gelidium microdon* Kützinger). BHT, DPPH and the methanol HPLC (high performance liquid chromatography) grade solvent were obtained from Sigma-Aldrich Chemical Co. (St. Louis, MO, USA). All chemicals used were of analytical grade. Deionized water used for the samples preparation was obtained with an in-house Milli-Q water purification system (Millipore, Bedford, MA, USA).

SEAWEEDS SAMPLE PREPARATION

The dried seaweed samples were cut into small pieces and ground into a fine powder using a dry grinder. The ground samples were sieved to obtain a uniform particle size and stored at -20 °C until further analysis. Each ground sample was weighed and transferred into a beaker. Methanol was added in the ratio of 1:10 and stirred for 2 hr with the aid of a magnetic stirrer. The extraction mixture was left to sediment for about 60 min before separating the extract from the residue by filtration through a cellulose acetate filter of 0.45 µm porosity. The residue was re-extracted twice with methanol in the same conditions and extracts were combined. The residual solvent was removed under reduced pressure at 45 °C using a

rotary evaporator (BÜCHI Labortechnik AG in Flawil, Switzerland). Extracts were prepared in duplicates and used to determine the FRSA.

FRSA ASSAY

The FRSA of the seaweeds species were determined in a methanolic solution of DPPH, used as a stable radical, according to the method described by Molyneux (2004) and Rainha et al. (2011a). This methodology measures the hydrogen atom or electron donor capacity of an extract to the stable radical DPPH formed in solution. In other words, it measures the capacity of the extract to scavenge free radicals in solution.

An aliquot of 2.0 mL of the methanolic solution of seaweed species or BHT at 2.0 mg.mL⁻¹ concentration was added to a test tube, with 1 mL of DPPH methanolic solution (4.5 mg.100 mL⁻¹ in methanol, w/v). Methanol was used as a blank to zero the absorbance (Abs), BHT was used as reference sample and a mixture without seaweed extract or BHT was used as the control (c). The Abs was measured at 517 nm over a period of 30 min and after each 5 min of reaction time using a Shimadzu 160-A UV/VIS spectrophotometer. All determinations were performed in triplicate and averaged. The FRSA of the samples (s) were calculated as a percentage of DPPH decolouration using the following equation (Rainha et al. 2011a):

$$\text{FRSA (\%)} = (1 - \text{Abs}_s/\text{Abs}_c) \times 100$$

Results are expressed as mean values \pm SD of three different Abs measurements of two extracts per seaweed species.

RESULTS

The antioxidant activity of the Azorean macroalgae compounds is mainly due to their redox properties, which can play an important role in adsorbing and neutralising free radicals, quenching singlet and triplet oxygen, or decomposing peroxides. Figure 1 shows the FRSA of the selected seaweed species that ranged from 19.54 \pm 2.40% to 60.05 \pm 4.29% after 30 minutes reaction. The SD of FRSA lower than 6.80% reveals good repeat-

ability. The FRSA results of all seaweed samples show slightly increased values with increasing reaction time, particularly from 10 to 30 min (Table 1). Among the methanolic seaweed extracts, the greatest anti-free radical activity after 30 minutes reaction was observed for *F. spiralis* (60.05 \pm 4.29%) followed by *G. microdon* (45.86 \pm 5.05%), *C. pachynema* (43.52 \pm 4.79%), *U. compressa* (39.12 \pm 4.30%), *O. pinnatifida* (33.97 \pm 3.74%), *P. capillacea* (27.44 \pm 5.90%), *Porphyra* sp. (22.73 \pm 4.88%) and *S. coronopifolius* (19.54 \pm 2.40%). The differences in FRSA values for different species reflect their different chemical compositions, probably due to differences in the type and amount of phenolic compound contents. However, the differences for the same species from different origins also reflect the influence of geographic origin, climate, season, variety (Dawes et al. 1998; Jiménez-Escrig & Cambrondón 1999) and seawater quality.




Fig. 1. Comparison of free radical scavenging activity (FRSA) of methanolic extract from selected Azorean macroalgae with standard butylated hydroxytoluene (BHT), at 2.0 mg.mL⁻¹ concentration, after 30 min reaction.

Table 1. Determination of free radical scavenging activity of selected Azorean macroalgae (reaction time from 5 min to 30 min)^a

Samples	Free radical scavenging activity (FRSA) % ^a					
	5 min	10 min	15 min	20 min	25 min	30 min
<i>Sphaerococcus coronopifolius</i>	7.05±3.41	10.72±5.20	13.50±1.84	15.82±2.16	17.69±1.54	19.54±2.40
<i>Porphyra</i> sp.	7.70±3.73	11.59±5.62	14.33±1.96	16.74±1.46	18.94±2.32	22.73±4.88
<i>Pterocladiaella capillacea</i>	17.34±1.51	19.19±2.35	21.33±3.27	22.80±4.90	25.56±5.49	27.44±5.90
<i>Osmundea pinnatifida</i>	20.50±3.14	25.86±5.56	27.97±6.01	29.57±6.35	31.50±3.47	33.97±3.74
<i>Ulva compressa</i>	16.15±1.41	23.04±4.95	29.15±6.26	32.86±3.62	35.98±3.96	39.12±4.30
<i>Chaetomorpha pachynema</i>	28.03±6.02	33.97±3.74	36.73±4.04	39.67±4.36	41.59±4.58	43.52±4.79
<i>Gelidium microdon</i>	30.44±6.54	34.90±3.84	38.68±4.26	41.40±4.55	44.31±4.88	45.86±5.05
<i>Fucus spiralis</i>	38.03±4.18	46.18±5.75	50.34±6.37	54.46±6.80	58.24±4.16	60.05±4.29
BHT	59.67±4.26	72.45±5.97	78.98±5.27	85.45±2.66	91.38±3.07	94.22±1.88

^aValues are expressed as mean ± SD (n = 6). Three determinations from two different batches of seaweeds were performed; BHT, butylated hydroxytoluene.

DISCUSSION

Generally, there is a preference for antioxidants from natural rather than from synthetic sources. The DPPH methodology allows one to determine the FRSA of different seaweeds with good repeatability. The results showed different values for different species, and *F. spiralis* and *G. microdon* presented superior antioxidant activities, revealing them to be valuable species that could be explored from biotechnology and commercial perspectives, taking into account the low seawater pollution levels in the Azores. The next step of this study will be the polyphenolic determination in order to establish the relationship with the antioxidant activity and also a fully antioxidant characterisation using other methodologies. These include: the β -carotene/linoleic acid assay (Dapkevicius et al. 1998), the reducing power effect assay (Oyaizu 1986), the superoxide anion scavenging activity assay (Liu et al. 1997), the ferrous ion-chelating assay (Dinis et al. 1994) and the bovine serum albumin (BSA) oxidative damage assay (Makris & Rossiter 2001), that were recently published with some experimental modifications by Rainha et al. (2011b).

ACKNOWLEDGEMENTS

The authors express their deep gratitude to the Biology Department technicians that collected the seaweeds. This work was supported by the grant

of CIRN and funds from Department of Technological Science and Development – University of Azores.

REFERENCES

- Conforti, F., M.R. Loizzo, G.A. Statti & F. Menichini 2005. Comparative radical scavenging and antidiabetic activities of methanolic extract and fractions from *Achillea ligustica* ALL. *Biological & Pharmaceutical Bulletin* 28: 1791–1794.
- Dapkevicius, A., R. Venskutonis, T.A. Van Beek & J. Linssen 1998. Antioxidant activity of extracts obtained by different isolation procedures from some aromatic herbs grown in Lithuania. *Journal of the Science of Food and Agriculture* 77: 140–146.
- Dastmalchi, K., H.J.D. Dorman, M. Kosar & R. Hiltunen 2007. Chemical composition and *in vitro* antioxidant evaluation of a water-soluble Moldavian blam (*Dracocephalum moldavica* L.) extract. *LWT - Food Science and Technology* 40: 239–248.
- Dawes, C.J. 1998. *Marine Botany*. John Wiley & Sons, Inc., New York. 480 pp.
- Dias, P.F., J.M. Siqueira Jr, L.F. Vendruscolo, T. DE Jesus Neiva, A.R. Gagliardi, M. Maraschin & R.M. Ribeiro-do-Valle 2005. Antiangiogenic and antitumoral properties of a polysaccharide isolated from the seaweed *Sargassum stenophyllum*. *Cancer Chemotherapy and Pharmacology* 56: 436–446.
- Dinis, T., V. Madeira & L. Alameida 1994. Action of phenolic derivatives (acetaminophen, salicylate, and 5-amino salicylate) as inhibitors of membrane

- lipid peroxidation and as peroxy radical scavengers. *Archives of Biochemistry and Biophysics* 315: 161–169.
- Duan, X.J., W.W. Zhang, X.M. Li & B.G. Wang 2006. Evaluation of antioxidant property of extract and fractions obtained from a red alga, *Polysiphonia urceolata*. *Food Chemistry* 95: 37–43.
- Duh, P.-D. 1998. Antioxidant activity of burdock (*Arctium lappa* Linné): Its scavenging effect on free radical and active oxygen. *Journal of the American Oil Chemists' Society* 75: 455–461.
- Ganesan, P., C.S. Kumar & N. Bhaskar 2008. Antioxidant properties of methanol extract and its solvent fractions obtained from selected Indian red seaweeds. *Bioresource Technology* 99: 2717–2723.
- Halliwell, B. 1991. Reactive oxygen species in living systems: source, biochemistry, and role in human disease. *American Journal of Medicine* 91: S14–S22.
- Jiménez-Escrig, A. & I.G. Cambrondón 1999. Evaluación nutricional y efectos fisiológicos de macroalgas marinas comestibles. *Archivos Latinoamericanos de Nutrición* 49: 114–120. [In Spanish]
- Jiménez-Escrig, A., I. Jiménez-Jiménez, R. Pulido & F. Saura-Calixto 2001. Antioxidant activity of fresh and processed edible seaweeds. *Journal of the Science of Food and Agriculture* 81: 530–534.
- Kuda, T., M. Tsunekawa, T. Hishi & Y. Araki 2005. Antioxidant properties of dried 'kayamo-nori', a brown alga *Scytosiphon lomentaria* (Scytosiphonales, Phaeophyceae). *Food Chemistry* 89: 617–622.
- Liu, F., V. Ooi & S. Chang 1997. Free radical scavenging activities of mushroom polysaccharide extracts. *Life Science* 60: 763.
- Makris, D. & J. Rossiter 2001. Comparison of quercetin and a non-orthohydroxy flavonol as antioxidants by competing in vitro oxidation reactions. *Journal of Agricultural and Food Chemistry* 49: 3370–3377.
- Matsukawa, R., Z. Dubinsky, E. Kishimoto, K. Masaki, Y. Masuda, T. Takeuchi, M. Chihara, Y. Yamamoto, E. Niki & I. Karube 1997. A comparison of screening methods for antioxidant activity in seaweeds. *Journal of Applied Phycology* 9: 29–35.
- Molyneux, P. 2004. The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin Journal of Science and Technology* 26: 211–219.
- Neto, A.I., V. Brotas, J.M.N. Azevedo, R.F. Patarra, N.M.V. Álavaro, C. Gameiro, A.C.L. Prestes & E.M. Nogueira 2009. *Qualidade de águas costeiras do Grupo Oriental do arquipélago dos Açores e proposta de monitorização*. Departamento de Biologia, Universidade dos Açores. iii+70 pp. [In Portuguese]
- Ohkawa, H., N. Ohishi & K. Yagi 1979. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical Biochemistry* 95: 351–358.
- Oyaizu, M. 1986. Studies on products of browning reactions: Antioxidative activities of products of browning reaction prepared from glucosamine. *Japanese Journal of Nutrition* 44: 307–315.
- Park, P.J., S.J. Heo, E.J. Park, S.K. Kim, H.G. Byun, B.T. Jeon & Y.J. Jeon 2005. Reactive oxygen scavenging effect of enzymatic extracts from *Sargassum thunbergii*. *Journal of Agricultural and Food Chemistry* 53: 6666–6672.
- Patarra, R.F., L. Paiva, A.I. Neto, E. Lima & J. Baptista 2011. Nutritional value of selected macroalgae. *Journal of Applied Phycology* 23(2): 205–208.
- Rackova, L., M. Oblozinsky, D. Kostalova, V. Kettman & L. Bezakova 2007. Free radical scavenging activity and lipoxygenase inhibition of Mahonia aquifolium extract and isoquinoline alkaloids. *Journal of Inflammation* 4: 15.
- Rainha, N., E. Lima & J. Baptista 2011a. Comparison of the endemic Azorean *Hypericum foliosum* with other *Hypericum* species. Antioxidant activity and phenolic profile. *Natural Product Research* 25(2): 123–135.
- Rainha, N., E. Lima, J. Baptista & C. Rodrigues 2011b. Antioxidant properties, total phenolic, carotenoid and chlorophyll content of anatomical parts of *Hypericum foliosum*. *Journal of Medicinal Plants Research* 5(10): 1930–1940.
- Rekka, E. & P.N. Kourounakis 1991. Effect of hydroxyethyl rutenosides and related compounds on lipid peroxidation and free radical scavenging activity. Some structural aspects. *Journal of Pharmacy and Pharmacology* 43: 486–491.
- Santoso, J., Y. Yoshie-Stark & T. Suzuki 2004. Antioxidant activity of methanol extracts from Indonesian seaweeds in an oil emulsion model. *Fisheries Science* 70: 183–188.
- Senevirathne, M., S.-H. Kim, N. Siriwardhana, J.-H. HA, K.-W. Lee & Y.-J. Jeon 2006. Antioxidant potential of *Ecklonia cava* on reactive oxygen species scavenging, metal chelating, reducing power and lipid peroxidation inhibition. *Food Science and Technology International* 12: 27–38.
- Siriwardhana, N., K.-W. Lee, Y.-J. Jeon, S.-H. Kim & J.-W. Haw 2003. Antioxidant activity of *Hizikia fusiformis* on reactive oxygen species scavenging and lipid peroxidation inhibition. *Food Science and Technology International* 9: 339–346.
- Spencer, J.P., A. Jenner, O.I. Aruoma, P.J. Evans, H. Kaur, D.T. Dexter, P. Jenner, A.J. Lees, D.C. Marsden & B. Halliwell 1994. Intense oxidative

- DNA damage promoted by L-DOPA and its metabolites. Implications for neurodegenerative disease. *FEBS Letters* 353: 246–250.
- Villaño, D., M.S. Fernández-Pachón, M.L. Moyá, A.M. Troncoso & M.C. Garcia-Parrilla 2007. Radical scavenging ability of polyphenolic compounds towards DPPH free radical. *Talanta* 71: 230–235.
- Yoshie, Y., W. Wang, Y.P. Hsieh & T. Suzuki 2002. Compositional difference of phenolic compounds between two seaweeds, *Halimeda* spp. *Journal of the Tokyo University of Fisheries* 88: 21–24.
- Received 25 Nov 2010. Accepted 2 Nov 2011, Published online 2 December 2011.*